
Azraq Jordan Refugee population 20,854 Total refugee population: Demographic breakdown: > 60 18% 21% 18 - 59 19% 17% 5 - 17 11% 10% 0 - 4



Indicators at a glance		
1 Health utilisation rate	8.3	X
2 Number of consultations per clinician per day	45	\checkmark
3 Hospitalisation rate (/1000/year)	61.0	
4 Referral rate (/1000/month)	16.1	
5 Crude Mortality Rate (CMR) (/1000/month)	0.3	\checkmark
6 Under-five Mortality Rate (U5MR) (/1000/month)	0.8	\checkmark
7 Neonatal Mortality Rate (NNMR) (/1000 livebirths)	14.0	\checkmark

%

Female

Notes


This public health profile for Azraq refugee camp is compiled from health facility data from the following partners:

• IMC Clinics
• IMC Hospital

Data is collected using standardised Health Information System (HIS) data collection and reporting tools and aims to be as accurate as possible.

Morbidity

Acute health conditions (%)



Top 5 acute health conditions


	% morbidity	# of cases
1 Upper Respiratory Tract Infection (URTI)	38%	25,813
2 Watery diarrhoea*	8%	5,050
3 Skin infection	7%	4,407
4 Influenza-like illness (ILI)	6%	3,823
5 Dental Conditions	5%	3,037
- Other	37%	25,037

Breakdown of acute health conditions

by age		by gender		
3% 38% 26%	■ 0-4 ■ 5 to 17 ■ 18 to 59 ■ > 60	50%	50%	■ Male ■ Female

UNHCR gratefully acknowledges the support of the following partners:


Chronic health conditions (%)



Top 5 chronic health conditions

	% morbidity	# of cases
1 Hypertension	38%	1,961
2 Diabetes	24%	1,206
3 Asthma	12%	587
4 Ischaemic heart disease	3%	178
5 Cardiovascular disorders (other)	3%	151
- Other	20%	1,012

Breakdown of chronic health conditions



Public Health			
Health Impact	Indicator	Standard	
Crude Mortality Rate (CMR) (/1000/month)	0.3	< 0.51	V
Under-five Mortality Rate (U5MR) (/1000/month)	0.8	< 1.5	V
Neonatal Mortality Rate (NNMR) (/1000 livebirths)	14.0	< 14.9	
Access and utilization			
No. of health facilities	1:6952	1:<10,000	V
Total number of consultations	86,461		
Health utilisation rate	8.3	1 - 4	X
No. of consultations per trained clinician per day	45	< 50	

Indicator Standard
0
100%
5,953

Breakdown of mental health conditions

By cause By gender By age ■ Epilepsy / seizures 18% Severe emotional disorder, including moderate-severe 32% 0-4 depression Intellecutal disability ■Male 48% ■5 to 17 ■ 18 to 59 ■ Female Psychotic disorder (including mania) > 60 28% ■ Other

Breakdown of injuries

By cause

By age

By gender

12%
5%
1%
Injury (burns)
Injury (war-related)
Injury (war-relate

	Indicator
Overall case fatality rate	0%
Proportion of deaths within 24 hours of admission	0%

Reproductive Health			
Antenatal care	Indicator	Standard	
Coverage of antenatal care (4 or more visits)	41%	≥ 90%	X
Coverage of TT vaccination	30%	≥ 90%	X
Coverage of anaemia screening in pregnancy	99%	≥ 90%	
Delivery care			
Proportion of deliveries attended by skilled personnel	100%	≥ 90%	
Proportion of deliveries performed by caesarean section	21%	5 - 15%	X
Proportion of low birth weight deliveries (<2500g)	2%	< 15%	
Maternal mortality			
Number of maternal deaths reported	0		
% of maternal deaths investigated within 48 hours	0%	100%	

Postnatal care	Indicator	Standard	
Coverage of postnatal care	36%	≥ 90%	X

Sexual and Gender-based Violence			
Incidence of rape (reported cases/10,000/year)*		^	
% of rape survivors who received PEP** < 72 hours	100%	^	
% of rape survivors who received ECP*** < 120 hours	100%	^	
% of rape survivors who received STI**** prophylaxis < 2 wks	100%	^	

*Only includes rapes reported at health facilities **Post-exposure prophylaxis ***Emergency contraceptive pill ****Sexually transmitted infection



















