

Camps reporting: Kawergosk, Qushtapa, Basirma and Darashakran

1. Population

| Population | 0 to < 1 | | Total < 1 | 1 to 4 | | Total 1 to 4 | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total |
|---------------------------------------|----------|-----|-----------|--------|-------|--------------|---------|-------|---------------|----------|-------|----------------|-------|-------|------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | M | F | | |
| Refugee population of reporting camps | 510 | 500 | 1,010 | 2,040 | 1,998 | 4,038 | 3,620 | 3,305 | 6,925 | 6,567 | 6,188 | 12,755 | 1,918 | 1,929 | 3,847 | 28,575 |
| % by age group | | | 4% | | | 14% | | | 24% | | | 45% | | | 13% | 100% |

* estimates from registered population in Kawergosk, Qushtapa, Basirma and Darashakran camps. For multiweek reports, mean is presented

2. Mortality

2.1 Mortality

| Record ONE cause of mortality per case | 0 to 4 | | Total < 5 | U5 % mort | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | Crude % mort |
|---|----------|----------|-----------|-------------|----------|----------|---------------|----------|----------|----------------|----------|----------|------------|-------------|--------------|
| | M | F | | | M | F | | M | F | | M | F | | | |
| 1. Lower Respiratory Tract Infection (LRTI) | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 2. Watery diarrhoea | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 3. Bloody diarrhoea | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 4. Cardiovascular disorders | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 5. Ischaemic heart disease | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 6. Injury | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 7. Cancer | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 8. Chronic obstructive pulmonary disease (COPD) | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 9. Tuberculosis (confirmed) | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 10. Neonatal death | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Other | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Total | 0 | 0 | 0 | 100% | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 100% |
| % by age group | | | | | | | | | | | | | | | |

| Mortality indicators | Male | Female | Total |
|--------------------------------|------|--------|-------|
| Crude Mortality Rate (CMR) | 0.0 | 0.0 | 0.0 |
| Under 5 Mortality Rate (U5MR) | 0.0 | 0.0 | 0.0 |
| Neonatal Mortality Rate (NNMR) | | | 0.0 |

2.2 Maternal death

| | |
|--|---|
| Number of maternal deaths reported | 0 |
| Number of maternal deaths investigated | 0 |
| Proportion of maternal deaths investigated | |

3. Morbidity

3.1 Staffing and service availability

| | |
|---|--------|
| Average number of full-time clinicians | 13.2 |
| Average number of full days on which each health facility was | 18.4 |
| Total number of outpatient consultations | 12,999 |
| Health facility utilisation rate | 5.9 |
| Consultations per clinician per day | 54 |

3.2 Acute health conditions

| Record NEW cases of disease only | 0 to 4 | | Total < 5 | U5 Incid** | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | % by cause | Crude Incid** |
|--|-------------|-------------|-------------|--------------|-------------|-------------|---------------|-------------|-------------|----------------|------------|------------|------------|--------------|-------------|---------------|
| | M | F | | | M | F | | M | F | | M | F | | | | |
| Upper Respiratory Tract Infection (URTI) | 1042 | 851 | 1893 | 93.8 | 744 | 660 | 1404 | 696 | 775 | 1471 | 46 | 36 | 82 | 4850 | 40% | 42.4 |
| Lower Respiratory Tract Infection (LRTI) | 257 | 178 | 435 | 21.5 | 107 | 86 | 193 | 103 | 103 | 206 | 21 | 13 | 34 | 868 | 7% | 7.6 |
| Watery diarrhoea* | 232 | 181 | 413 | 20.5 | 34 | 37 | 71 | 32 | 27 | 59 | 7 | 4 | 11 | 554 | 5% | 4.8 |
| Bloody diarrhoea* | 6 | 9 | 15 | 0.7 | 0 | 0 | 0 | 3 | 0 | 3 | 0 | 0 | 0 | 18 | 0% | 0.2 |
| Malaria (suspected)* | 2 | 1 | 3 | 0.1 | 3 | 1 | 4 | 4 | 2 | 6 | 0 | 0 | 0 | 13 | 0% | 0.1 |
| Skin infection | 62 | 63 | 125 | 6.2 | 51 | 46 | 97 | 107 | 134 | 241 | 8 | 7 | 15 | 478 | 4% | 4.2 |
| Eye infection | 35 | 38 | 73 | 3.6 | 53 | 18 | 71 | 74 | 72 | 146 | 9 | 8 | 17 | 307 | 3% | 2.7 |
| Ear infection | 77 | 75 | 152 | 7.5 | 46 | 54 | 100 | 56 | 103 | 159 | 10 | 7 | 17 | 428 | 4% | 3.7 |
| Dental Conditions | 12 | 12 | 24 | 1.2 | 34 | 33 | 67 | 125 | 176 | 301 | 17 | 12 | 29 | 421 | 4% | 3.7 |
| Intestinal Worms | 7 | 18 | 25 | 1.2 | 18 | 13 | 31 | 7 | 6 | 13 | 1 | 0 | 1 | 70 | 1% | 0.6 |
| Acute Jaundice Syndrome* | 0 | 1 | 1 | 0.0 | 0 | 1 | 1 | 1 | 0 | 1 | 0 | 0 | 0 | 3 | 0% | 0.0 |
| Measles (suspected)* | 0 | 1 | 1 | 0.0 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0% | 0.0 |
| Meningitis (suspected)* | 0 | 1 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0% | 0.0 |
| Acute Flaccid Paralysis* | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Tuberculosis (suspected) | 1 | 0 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0% | 0.0 |
| Fever of unknown origin | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Sexually Transmitted Infection (STI) | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0% | 0.0 |
| Urinary tract infection (UTI) | 18 | 41 | 59 | 2.9 | 30 | 51 | 81 | 115 | 361 | 476 | 44 | 9 | 53 | 669 | 6% | 5.9 |
| HIV/AIDS | 0 | 0 | 0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% | 0.0 |
| Acute malnutrition | 2 | 0 | 2 | 0.1 | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 | 4 | 0% | 0.0 |
| Leishmaniasis | 0 | 1 | 1 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0% | 0.0 |
| Other | 218 | 150 | 368 | 18.2 | 197 | 151 | 348 | 838 | 1443 | 2281 | 172 | 161 | 333 | 3330 | 28% | 29.1 |
| Total | 1971 | 1621 | 3592 | 177.9 | 1318 | 1152 | 2470 | 2163 | 3202 | 5365 | 335 | 257 | 592 | 12019 | 100% | 105.2 |
| % by age group | | | 30% | | | | 21% | | | 45% | | | 5% | 100% | | |

* indicates disease with epidemic potential. See guidance for alert thresholds and advice on how these should be interpreted. **incidence is cases per 1000 per week

3.3 Chronic health conditions

| Record NEW and REVISITS | 0 to 4 | | Total < 5 | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | % by cause | |
|--|-----------|-----------|-----------|-----------|-----------|---------------|-----------|------------|----------------|------------|------------|------------|-------------|-------------|--|
| | M | F | | M | F | | M | F | | M | F | | | | |
| Diabetes | 35 | 9 | 44 | 14 | 22 | 36 | 24 | 40 | 64 | 24 | 77 | 101 | 245 | 35% | |
| Hypertension | 4 | 3 | 7 | 0 | 4 | 4 | 21 | 98 | 119 | 47 | 129 | 176 | 306 | 44% | |
| Asthma | 5 | 5 | 10 | 1 | 3 | 4 | 7 | 4 | 11 | 13 | 1 | 14 | 39 | 6% | |
| Ischaemic heart disease | 0 | 0 | 0 | 0 | 0 | 0 | 7 | 9 | 16 | 10 | 11 | 21 | 37 | 5% | |
| Cardiovascular disease (other) | 0 | 0 | 0 | 1 | 1 | 2 | 5 | 13 | 18 | 4 | 7 | 11 | 31 | 4% | |
| Chronic obstructive pulmonary disease (COPD) | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 2 | 0 | 2 | 3 | 0% | |
| Liver disease | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 3 | 0 | 1 | 1 | 4 | 1% | |
| Thyroid problems | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 10 | 12 | 0 | 4 | 4 | 16 | 2% | |
| Cancer | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 6 | 7 | 1 | 0 | 1 | 8 | 1% | |
| Other | 0 | 1 | 1 | 0 | 1 | 1 | 4 | 4 | 8 | 2 | 0 | 2 | 12 | 2% | |
| Total | 44 | 18 | 62 | 16 | 31 | 47 | 72 | 187 | 259 | 103 | 230 | 333 | 701 | 100% | |
| % by age group | | | 9% | | | 7% | | | 37% | | | 48% | 100% | | |

3.4 Mental health

| | 0 to 4 | | Total < 5 | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | % by cause |
|---|----------|----------|-----------|----------|----------|---------------|-----------|-----------|----------------|----------|----------|------------|-------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| 1. Epilepsy / seizures | 0 | 1 | 1 | 2 | 1 | 3 | 7 | 8 | 15 | 0 | 0 | 0 | 19 | 22% |
| 2. Alcohol and other substance abuse disorder | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0% |
| 3. Psychotic disorder | 0 | 0 | 0 | 0 | 0 | 0 | 9 | 0 | 9 | 0 | 0 | 0 | 9 | 11% |
| 4. Mental retardation / intellectual disability | 0 | 0 | 0 | 0 | 2 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 2% |
| 5. Severe emotional disorder | 0 | 1 | 1 | 0 | 1 | 1 | 2 | 5 | 7 | 0 | 0 | 0 | 9 | 11% |
| 6. Medically unexplained somatic complaint | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 2 | 0 | 1 | 1 | 3 | 4% |
| Other | 4 | 0 | 4 | 2 | 2 | 4 | 7 | 24 | 31 | 1 | 3 | 4 | 43 | 51% |
| Total | 4 | 2 | 6 | 4 | 6 | 10 | 25 | 39 | 64 | 1 | 4 | 5 | 85 | 100% |
| % by age group | | | 7% | | | 12% | | | 75% | | | 6% | 100% | |

3.5 Injuries

| | 0 to 4 | | Total < 5 | 5 to 14 | | Total 5 to 14 | 15 to 49 | | Total 15 to 49 | ≥ 50 | | Total ≥ 50 | Grand total | % by cause |
|---------------------------|-----------|-----------|-----------|-----------|-----------|---------------|-----------|-----------|----------------|----------|----------|------------|-------------|-------------|
| | M | F | | M | F | | M | F | | M | F | | | |
| 1. Injury (intentional) | 0 | 0 | 0 | 2 | 1 | 3 | 4 | 0 | 4 | 0 | 0 | 0 | 7 | 4% |
| 2. Injury (unintentional) | 12 | 8 | 20 | 19 | 14 | 33 | 52 | 7 | 59 | 4 | 0 | 4 | 116 | 60% |
| 3. Burns | 5 | 3 | 8 | 4 | 4 | 8 | 9 | 9 | 18 | 0 | 2 | 2 | 36 | 19% |
| 4. Accident | 1 | 1 | 2 | 2 | 0 | 2 | 10 | 2 | 12 | 1 | 0 | 1 | 17 | 9% |
| Other | 2 | 1 | 3 | 3 | 2 | 5 | 7 | 3 | 10 | 0 | 0 | 0 | 18 | 9% |
| Total | 20 | 13 | 33 | 30 | 21 | 51 | 82 | 21 | 103 | 5 | 2 | 7 | 194 | 100% |
| % by age group | | | 17% | | | 26% | | | 53% | | | 4% | 100% | |

4. Referrals

4.1 Referrals - general

| | 0 to 4 | 5 to 14 | 15 to 49 | ≥ 50 | Grand total |
|----------------|-----------|-----------|------------|-----------|-------------|
| Emergency | 5 | 8 | 21 | 4 | 38 |
| Elective | 25 | 23 | 106 | 46 | 200 |
| Other | 15 | 7 | 34 | 2 | 58 |
| Total | 45 | 38 | 161 | 52 | 296 |
| % by age group | 15% | 13% | 54% | 18% | 100% |

5. Laboratory & other investigations

| | Total |
|-------------------------|-----------|
| Haematology | 0 |
| Imaging (X-ray, MRI/CT) | 0 |
| Other | 0 |
| Total | 12 |

6. EPI and Vitamin A

| | <1 | ≥1 to <5 | Total |
|--|----|----------|-------|
| BCG | 0 | 0 | 0 |
| Hepatitis B birth dose | 0 | 0 | 0 |
| DPT I/HiB I/Hep B I* (Pentavalent I) | 0 | 0 | 0 |
| DPT II/HiB II* (Tetravalent) | 0 | 0 | 0 |
| DPT III/HiB III/Hep B II* (Pentavalent II) | 0 | 0 | 0 |
| Polio 0 | 0 | 0 | 0 |
| Polio I | 0 | 0 | 0 |
| Polio II | 0 | 0 | 0 |
| Polio III | 0 | 0 | 0 |
| Measles | 0 | 0 | 0 |
| MMR | 0 | 0 | 0 |
| Rotavirus 1 | 0 | 0 | 0 |
| Rotavirus 2 | 0 | 0 | 0 |
| Rotavirus 3 | 0 | 0 | 0 |
| Vitamin A | 0 | 0 | 0 |
| Additional vaccines | 0 | 4 | 4 |

| EPI Indicators (since December 2013) | | |
|--------------------------------------|--|------|
| Total vaccinated | | 4 |
| a. Program dropout rate* | | -30% |
| b. Immunisation coverage | | |
| BCG | | 63% |
| Hepatitis B birth dose | | 50% |
| DPT I | | 69% |
| DPT III | | 88% |
| Polio I | | 72% |
| Polio III | | 91% |
| Measles | | 116% |
| MMR** | | 111% |
| Rotavirus I | | 60% |
| Rotavirus III | | 76% |
| c. Vitamin A coverage | | 103% |

*If program dropout <0%, please review data recording and reporting procedures for multi-dose vaccines
Targets: Program dropout, 0 to <10%. Immunisation coverage, ≥90%
**For most indicators, coverage estimates calculated for <1 years. MMR calculated for 1 to <2 years.

Note: If actual population living in the camps is lower than reported population, coverage indicators will be artificially depressed.

7. Reproductive Health

7.1 Antenatal care

| | < 18 | ≥ 18 | Total |
|--|----------------|------|-------|
| First antenatal visit < 1st trimester | | | |
| First antenatal visit > 1st trimester | | | |
| Number of pregnant women at time of delivery who received: | NO DATA | | |
| ≥ 4 ANC visits | | | |
| 2 TT doses | | | |
| anaemia screening | | | |

ANC Indicators

| | |
|--|--|
| Proportion of first time antenatal visits made < 1st trimester | |
|--|--|

7.2 Postnatal care

| | < 18 | ≥ 18 | Total |
|--|------|------|----------------|
| Women who attended 3 postnatal visits within 6 wks of delivery | | | NO DATA |

7.3 Sexual and gender-based violence

| | < 18 | | ≥ 18 | | Total |
|---|------|---|------|---|-------|
| | M | F | M | F | |
| Total no rape survivors seen | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 72 -120 hours | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined within 120 hrs - 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. of rape survivors examined after 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No of rape survivors given PEP within 72 hours | 0 | 0 | 0 | 0 | 0 |
| No of female rape survivors given ECP within 120 hours | | | | | 0 |
| No of rape survivors given STI presumptive treatment within 2 weeks | 0 | 0 | 0 | 0 | 0 |
| No. cases of trauma in health centre due to domestic violence | 0 | 0 | 0 | 0 | 0 |

ECP = Emergency Contraceptive Pill; PEP = Post-exposure Prophylaxis; STI = Sexually Transmitted Infection

SGBV Indicators

| | |
|--|-----|
| Incidence of reported rape | 0.0 |
| Proportion of rape survivors who receive PEP < 72 hrs | |
| Proportion of female rape survivors who receive ECP < 120 hrs | |
| Proportion of rape survivors who receive STI presumptive treatment < 2 wks | |

Data Sources

1. Health Information System

Publication date:

15 May 2014

Contact Information

Erbil
Dr Raad Najmaddin, DOH Erbil,
Email: RAAD976@yahoo.com

Erbil
Sandra Harlass, UNHCR
Email: HARLASS@unhcr.org

We gratefully acknowledge the contributions of the following agencies:

