Azraq Camp | Jordan

Refugee population

Total refugee population: **37,157**

Demographic breakdown:

Age group	Male	Female
≥ 60	-1%	1%
18 - 59	-19%	22%
5 - 17	-19%	18%
0 - 4	-10%	10%

Male % Female

Key facts

Camp opened:
May 2014

Indicators at a glance

|			
1	Health utilisation rate	6.1	X
2	Number of consultations per clinician per day	35	√
3	Referral rate (/1,000/month)	6.3	
4	Crude Mortality Rate (CMR) (/1,000/month)	0.2	√
5	Under-five Mortality Rate (U5MR) (/1,000/month)	0.3	√
6	Neonatal Mortality Rate (NNMR) (/1,000 livebirths)	10.6	√

Notes

This public health profile for Azraq refugee camp is compiled from health facility data from the following partners:
- **IMC Clinics in Villages 3, 5, and 6**
- **IMC Hospital**
- **Arab Medical Relief Clinics in Villages 2 and 6**
- **IRC RH Services Villages 3, 5, and 6**

Data is collected using standardised Health Information System (HIS) data collection and reporting tools and aims to be as accurate as possible.

Morbidity

Acute health conditions (%)

- Upper Respiratory Tract Infection (URTI) — 21%
- Dental Conditions — 15%
- Influenza-like illness (ILI) — 5%
- Skin infection — 4%
- Watery diarrhoea* — 3%
- Other — 52%

Chronic health conditions (%)

- Hypertension — 34%
- Diabetes — 20%
- Asthma — 8%
- Thyroid problems — 6%
- Ischaemic heart disease — 5%
- Other — 27%

Top 5 acute health conditions

|		% morbidity	# of cases
1	Upper Respiratory Tract Infection (URTI)	21%	15,226
2	Dental Conditions	15%	10,788
3	Influenza-like illness (ILI)	5%	3,856
4	Skin infection	4%	2,848
5	Watery diarrhoea*	3%	2,127
-	Other	52%	38,022

Top 5 chronic health conditions

		% morbidity	# of cases
1	Hypertension	34%	2,058
2	Diabetes	20%	1,176
3	Asthma	8%	486
4	Thyroid problems	6%	376
5	Ischaemic heart disease	5%	282
-	Other	27%	1,650

Breakdown of acute health conditions

by age
- 0-4 — 28%
- 5 to 17 — 26%
- 18 to 59 — 43%
- > 60 — 3%

by gender
- Male — 44%
- Female — 56%

Breakdown of chronic health conditions

by age
- 0-4 — 4%
- 5 to 17 — 5%
- 18 to 59 — 66%
- > 60 — 25%

by gender
- Male — 44%
- Female — 56%

UNHCR gratefully acknowledges the support of the following partners:

Public Health

Health Impact

| Health Impact	Indicator	Standard	
Crude Mortality Rate (CMR) (/1,000/month)	0.2	< 0.5	√
Under-five Mortality Rate (U5MR) (/1,000/month)	0.3	< 1.5	√
Neonatal Mortality Rate (NNMR) (/1,000 livebirths)	10.6	< 14.9	√

Access and utilization

Access and utilization	Indicator	Standard	
No. of health facilities	1 : 7432	1 : < 10,000	√
Total number of consultations	112,058		
Health utilisation rate	6.1	1 - 4	X
No. of consultations per trained clinician per day	35	< 50	√

Breakdown of mental health conditions

By cause

- Severe emotional disorder, including moderate-severe depression — 44%
- Epilepsy / seizures — 21%
- Psychotic disorder (including mania) — 6%
- Intellecutal disability — 4%
- Other — 25%

By age

- 0-4 — 3%
- 5 to 17 — 29%
- 18 to 59 — 63%
- > 60 — 5%

By gender

- Male — 46%
- Female — 54%

Breakdown of injuries

By cause

- Injury (burns) — 27%
- Bites (all causes)
- Injury (war-related)
- Injury (war-related)
- Other — 73%

By age

- 0-4 — 23%
- 5 to 17 — 33%
- 18 to 59 — 41%
- > 60 — 3%

By gender

- Male — 50%
- Female — 50%

In-patient and referral

In-patient department	Indicator	Standard	
Hospitalisation rate (/1,000/year)	75.5	50 - 150	^
Bed occupancy rate	61%	> 75%	X
Mean length of stay for authorised discharges (days)	2.1		

	Indicator	Standard
Overall case fatality rate	0%	
Proportion of deaths within 24 hours of admission	0%	

Reproductive Health

Antenatal care	Indicator	Standard	
Coverage of antenatal care (4 or more visits)	73%	≥ 90%	X
Coverage of TT vaccination	13%	≥ 90%	X
Coverage of anaemia screening in pregnancy	99%	≥ 90%	√

Delivery care	Indicator	Standard	
Proportion of deliveries attended by skilled personnel	98%	≥ 90%	√
Proportion of deliveries performed by caesarean section	26%	5 - 15%	X
Proportion of low birth weight deliveries (<2500g)	7%	< 15%	√

Maternal mortality	Indicator	Standard
Number of maternal deaths reported	1	
% of maternal deaths investigated	100%	100%

Postnatal care	Indicator	Standard	
Coverage of postnatal care	33%	≥ 90%	X

Sexual and Gender-based Violence	Indicator	Standard		
Incidence of rape (reported cases/10,000/year)*	0.0		^	⚑
% of rape survivors who received PEP** < 72 hours		100%	^	⚑
% of rape survivors who received ECP*** < 120 hours		100%	^	⚑
% of rape survivors who received STI**** prophylaxis < 2 wks		100%	^	⚑

*Only includes rapes reported at health facilities **Post-exposure prophylaxis ***Emergency contraceptive pill ****Sexually transmitted infection